



IMEDEA



# NOISE-SUSTAINED PATTERNS IN NONLINEAR OPTICS

Pere Colet, Maxi San Miguel, Roberta Zambrini

Marco Santagiustina, [Università di Padova, Italy](#)

Gonzalo Izús, [Universidad de Mar del Plata, Argentina](#)

Daniel Walgraef, [Université Libre de Bruxelles, Belgium](#)

## 3 REGIMES

- **Absolutely Stable:**


Perturbation decays

- **Absolutely Unstable:**

Perturbation grows and spreads faster than the advection velocity

- **Convectively Unstable:**

Perturbation grows but spreads slower than the advection velocity



Absolute Instability threshold:  $v_F = v$

Convectively unstable regime  
+  
noise



Noise-sustained patterns

## Taylor-Couette flow




FIG. 7. Space-time plots of noise-sustained structure in the convectively unstable regime. (a) Experimental flow visualization with Kalliroscope for  $R=3.0$ ,  $\nu=0.025$  S, and  $\epsilon=0.040$ . (b) Integration of Eq. (5.2) with noise level  $F_A=2.1 \times 10^{-7}$ .

## Variance of time spectra at a fixed position




FIG. 10. Time series of reflected light, and their DFT moduli near the fundamental peak, at  $z=100$  for  $R=3.0$ . About  $\frac{1}{4}$  of each series is shown. (a)  $\epsilon - \epsilon_c = 0.0896$ . (b)  $\epsilon - \epsilon_c = 0.0318$ . (c) Structure persists at small  $\epsilon - \epsilon_c = 0.0077$ . The DFT peak shows considerable broadening.




FIG. 11. (a) Normalized second moment of the fundamental peak in the DFT power vs  $\epsilon$  at  $R=3.0$  for experimental time series at  $z=100$ . The transition at  $\epsilon=0.065$  indicates the onset of phase noise. (b) Corresponding results for Eq. (5.2) for inlet noise levels  $10^{-6}$  (+),  $10^{-5}$  ( $\Delta$ ), and  $10^{-4}$  ( $\circ$ ), respectively. The dashed line locates  $\epsilon_c^{\text{CGLE}}=0.071$ .

## **CONCEPTS IN NOISE SUSTAINED STRUCTURES**

***PHENOMENON OF NOISE AMPLIFICATION***, while noisy precursors are weakly damped critical fluctuations

***NOISE NEEDED AT ALL TIMES:*** A laser requires amplification of spontaneous emission, but when the laser is lasing noise is no longer required to maintain the oscillation

***PRECURSOR OF AN ABSOLUTELY UNSTABLE REGIME:***  
An optical amplifier is in a convective regime, but there is no regime of absolute instability


---

# Noise-sustained patterns in optical cavities


---

---

## Examples



# 1d self-focusing Kerr cavities



$$E(x, z, t) = \underbrace{A(x, t)}_{\text{field envelope}} e^{i(k_0 z - \omega_0 t)}$$

## Time evolution of field envelope

$$\partial_t A - 2\alpha_0 \partial_x A = -(1 + i\Delta)A + ia\partial_x^2 A + E_0 + i|A|^2 A + \sqrt{\epsilon}\xi(t)$$


$\alpha_0$ : tilt angle       $a$ : diffraction strength       $E_0$ : pump  
 $\Delta$ : cavity detuning       $\xi_i$ : Gaussian white noise

Homogeneous solution:  $E_0 = A^{\text{st}} \left[ 1 + i(\Delta - |A^{\text{st}}|^2) \right]$

Linear stability analysis:  $A(x, t) = A^{\text{st}} + \delta A(x, t)$

$$\delta A(x, t) = \sum_q b_q(t) e^{iqx} \quad b_q(t) \propto e^{\lambda(q)t}$$

$$\lambda_{\pm}(q) = 2i\alpha_0 q - 1 \pm \sqrt{|A^{\text{st}}|^4 - (2|A^{\text{st}}|^2 - \Delta - aq^2)^2}$$



Instability threshold:  $\text{Re}(\lambda_1) = 0 \Rightarrow$

$$\begin{cases} \Delta < 2: & |A|^2 = 1 & |q_c| = \sqrt{2 - \Delta}/a & \text{pattern} \\ \Delta > 2: & |A|^2 = \frac{2\Delta - \sqrt{\Delta^2 - 3}}{3} & |q_c| = 0 & \text{homogeneous} \end{cases}$$

# Convective stability analysis

Linearized asymptotic behaviour of a generic perturbation  $\psi$  of steady state:

$$\psi(x,t) = \int_{-\infty}^{+\infty} dq \tilde{\psi}(q,0) \exp[iqx + \lambda(q)t] \quad \begin{array}{l} \tilde{\psi}(q,0) : \text{initial perturbation in Fourier space} \\ \lambda(q) : \text{eigenvalue with largest real part} \end{array}$$

$\text{Re}[\lambda(q)] < 0, q \in R, \Rightarrow \psi(x,t) \rightarrow 0$  as  $t \rightarrow \infty$  **Absolutely stable**

$\text{Re}[\lambda(q)] > 0$   $\left\{ \begin{array}{l} |\psi(x,t)| \rightarrow \infty \text{ for arbitrary } x \\ |\psi(x_0,t)| \rightarrow 0 \text{ and } |\psi(x_0 + vt, t)| \rightarrow \infty \text{ for some } v \end{array} \right.$  **Absolutely unstable**

**Convectively unstable**

To evaluate the integral:

- Analytically continue  $\lambda(q)$  over complex plane  $k$  in the form  $\lambda(k/i)$
- Deform integration contour to pass through saddle point  $k^s$  along direction of steepest descent.

$$\partial_k \lambda(k)|_{k^s} = 0, \quad \text{Re} \left[ \partial_k^2 \lambda(k)|_{k^s} \right] \geq 0$$

The system is

**Absolutely unstable** for:  $\text{Re}[\lambda(q)] > 0, \text{Re}[\lambda(k^s)] > 0$

**Convectively unstable** for:  $\text{Re}[\lambda(q)] > 0, \text{Re}[\lambda(k^s)] < 0$


Deissler, J. Stat. Phys., **40**, 376 (85); **54**, 1459 (89); Physica D **56**, 303 (92);  
Hall & Heckrotte, Phys. Rev. **166**, 120 (68); T.H. Stix, *Waves in plasmas*, AIP (92);  
Infield & Rowlands, *Nonlinear waves, solitons and chaos*, Cambridge (90)

# 1d self-focusing Kerr cavities

Analytic continuation of  $\lambda_+(q)$  over complex plane:  $\lambda(k) = 2\alpha_0 k - 1 + \sqrt{|A^{st}|^4 - (2|A^{st}|^2 - \Delta + ak^2)^2}$


Saddle:  $\partial_k \lambda(k)|_{k^s} = 0 \Rightarrow \alpha_0 - \frac{2a(2|A^{st}|^2 - \Delta + ak^2)k^s}{\sqrt{|A^{st}|^4 - (2|A^{st}|^2 - \Delta + ak^2)^2}} = 0$

Absolute instability threshold:  $\text{Re}[\lambda(k^s)] = 0$






# 1d self-focusing Kerr cavities



Variance of time spectra at a fixed position




Spectral narrowing identifies transition from noise-sustained to deterministic pattern

# 2d self-focusing Kerr cavities

## Absolutely unstable:


Stability analysis shows first unstable modes are stripes perpendicular to tilt.

Stripes not stable in nonlinear regime hexagons.





## Convectively unstable:

Noise generates disordered hexagons preceded by unstable noisy stripes.



**NOISE SUSTAINED STRUCTURE:  
MACROSCOPIC AND SPATIALLY STRUCTURED  
MANIFESTATION OF QUANTUM NOISE**

# Type I degenerate optical parametric oscillator



pump( $2\omega$ ):  $A_0$   $\partial_t A_0 = \gamma_0 \left[ -(1+i\Delta_0)A_0 + E_0 + ia_0 \nabla^2 A_0 + 2iK_0 A_1^2 \right] + \sqrt{\epsilon_0} \xi_0(\vec{x}, t)$

signal( $\omega$ ):  $A_1$   $\partial_t A_1 = \gamma_1 \left[ -(1+i\Delta_1)A_1 + ia_1 \nabla^2 A_1 + iK_0 A_1^* A_0 + \rho \partial_y A_1 \right] + \sqrt{\epsilon_1} \xi_1(\vec{x}, t)$

$\gamma_i$ : cavity decay rates     $a_i$ : diffraction strength     $E_0$ : pump  
 $\Delta_i$ : cavity detunings     $\xi_i$ : Gaussian white noise     $K_0$ : nonlinear coef.

**Walk-off**  $\rightarrow$  **convection term**

Homogeneous solution:  $A_0^{\text{st}} = E_0 / (1 + i\Delta_0)$ ,  $A_1^{\text{st}} = 0$

$F \equiv K_0 |A_0^{\text{st}}|$

$A_j(\vec{x}, t) = A_j^{\text{st}} + \delta A_j(\vec{x}, t)$

$\delta A_j(\vec{x}, t) = \sum_{\vec{q}} b_{j\vec{q}}(t) e^{i\vec{q}\vec{x}}$      $b_{j\vec{q}}(t) \propto e^{\lambda_j(\vec{q})t}$

$$\begin{cases} \lambda_0(\vec{q}) = -\gamma_0(1 + i\Delta_0 + ia_0 q^2) & \text{Re}(\lambda_0) < 0 \\ \lambda_{1\pm}(\vec{q}) = \gamma_1 \left[ -1 + i\rho q_y \pm \sqrt{F^2 - (\Delta_1 + a_1 q^2)} \right] \end{cases}$$

Instability threshold:  $\text{Re}(\lambda_{1+}) = 0 \Rightarrow F^2 = 1$

$$\begin{cases} \Delta_1 < 0: & |\vec{q}_c| = \sqrt{-\Delta_1 / a_1} & \text{pattern} \\ \Delta_1 > 0: & |\vec{q}_c| = 0 & \text{homogeneous} \end{cases}$$


# Convective stability analysis

Analytic continuation of  $\lambda_{1+}(\vec{q})$  over complex plane:  $\lambda(\vec{k}) = \gamma_1 \left[ -1 + \rho k_y + \sqrt{F^2 - a_1(q_c^2 + k^2)^2} \right]$

Saddle:  $\nabla_{\vec{k}} \lambda(k_x, k_y)|_{k_x^s, k_y^s} = 0 \Rightarrow \begin{cases} k_x^s = 0 \Rightarrow q_x = 0 & \text{pattern orientation perpendicular to drift} \\ \rho - \frac{2a_1^2(q_c^2 + k_y^{s2})k_y^s}{\sqrt{F^2 - a_1(q_c^2 + k_y^{s2})^2}} = 0 \end{cases}$

Absolute instability threshold:  $\text{Re}[\lambda(0, k_y^s)] = 0$

- Abs. unstable
- Conv. unstable
- Abs. stable



M. Santagiustina, P. Colet, M. San Miguel and D. Walgraef, Phys. Rev. E **58**, 3843 (1998) ;

H. Ward, M.N. Ouarzazi, M. Taki, P. Glorieux, Eur. Phys. J. D **3**, 275 (1998);

M. Taki, M.N. Ouarzazi, H. Ward, P. Glorieux, J. Opt. Soc. Am. B **17**, 997 (2000).

# Type I degenerate optical parametric oscillator




---

absolutely stable




convectively unstable

absolutely unstable

$|A_1(x,y)|^2$




$|A_1(q_x, q_y)|^2$




*M. Santagiustina, P. Colet, M. San Miguel and D. Walgraef, PRE 58, 3843 (1998)*

---

# Type I degenerate optical parametric oscillator



Variations of spectra at different spatial positions.



Large spread due to relatively short integration time.




# Type I DOPO. Wavevector selection

Periodic array of dark solitons



Selected wavevector  $Q_y$ :

- The linearly unstable wavevector
  - oscillates at frequency  $\text{Im}[\lambda(\vec{k}^s)]$
  - drifts at velocity  $\rho$ .
- Imposing conservation of field node flux from linearly unstable regime to the stable one:

$$Q_y = \frac{\text{Im}[\lambda(\vec{k}^s)]}{\rho}$$



# Type II Optical Parametric Oscillator



**Type-II interaction**

Polarization plays a fundamental role.

pump( $2\omega$ ), ordinary:  $A_0$

signal( $\omega$ ), ordinary:  $A_1$

idler( $\omega$ ), extraordinary:  $A_2$

$$\partial_t A_0 = \gamma_0 \left[ -(1 + i\Delta_0) A_0 + E_0 + ia_0 \nabla^2 A_0 + 2iK_0 A_1 A_2 \right] + \sqrt{\epsilon_0} \xi_0(\mathbf{r}, t)$$

$$\partial_t A_1 = \gamma_1 \left[ -(1 + i\Delta_1) A_1 + ia_1 \nabla^2 A_1 + iK_0 A_2^* A_0 \right] + \sqrt{\epsilon_1} \xi_1(\mathbf{r}, t)$$

$$\partial_t A_2 = \gamma_2 \left[ -(1 + i\Delta_2) A_2 + ia_2 \nabla^2 A_2 + iK_0 A_1^* A_0 + \rho_2 \partial_y A_2 \right] + \sqrt{\epsilon_2} \xi_2(\mathbf{r}, t)$$

**Walk-off  $\rightarrow$  convection term**

$\gamma_i$ : cavity decay rates

$a_i$ : diffraction strength

$E_0$ : pump

$\Delta_i$ : cavity detunings

$\xi_i$ : Gaussian white noise

$K_0$ : nonlinear coef.



# Type II OPO: Instabilities of the solution $A_1=A_2=0$

- For small  $E_0$ ,  $A_{1,2}=0$  is a stable solution

- For  $\Delta_{1,2} < 0$ , zero solution unstable at:

$$F = |K_0 E_0 / (1 + i \Delta_0)| = 1.$$



Instability takes place at finite wavevector for  $A_1$  and  $A_2$

Close to threshold, instability is convective.

- Larger pump: absolute instability.

One transverse  $q$  mode, parallel to the walk-off direction, is selected for  $A_1$  and the opposite  $-q$  for  $A_2$ .

Two equally probable selections: (□, □) and (○, ○).




# Type II OPO: Absolutely unstable regime

## Transient regime

- Competition between two travelling waves of different wavevector (modes  $\bigcirc$  and  $\square$ )
- Intensity constant in each travelling wave domain.
- **defects appear** at domain boundaries


## Final state

- One travelling wave takes over.
- Selection is random.
- No intensity patterns, only phase stripes




# Type II OPO: Convectively unstable regime

$$|A_1(\mathbf{x}, \mathbf{y})|^2$$



X

$$|A_1(\mathbf{q}_x, \mathbf{q}_y)|^2$$



In the convective regime, all spatial modes are sustained by noise, partially restoring symmetry

Intensity stripes of random direction appear at bottom of pattern.



They orient themselves while they are amplified and drift

**Interference of the modes sustained by noise generates an intensity pattern**

# Type II OPO: Synchronization of noise-sustained patterns

$$|A_1(x,y)|^2$$

$$|A_2(x,y)|^2$$



Despite the fact that patterns in  $A_1$  and  $A_2$  are stochastic, macroscopic size structures driven by independent noises, their correlation grows with time and finally leads to synchronization of noise-sustained structures.

$$\Gamma(P,Q) = \frac{\iint (P - \langle P \rangle)(Q - \langle Q \rangle) dx dy}{\sqrt{\sigma(P)}\sqrt{\sigma(Q)}}$$

